



*This paper forms part of the proceedings from the BOU conference **Marine Renewables and Birds**  
Other papers from these proceedings can be viewed at [www.BOUPROC.net](http://www.BOUPROC.net).*

POSTER ABSTRACT

**Examining between-year natural variation in seabird numbers and distribution  
at an offshore wind farm development site**

**COLIN BARTON<sup>1\*</sup>, MARTIN POOT<sup>2</sup>, ZOË CRUTCHFIELD<sup>3</sup>, PHIL BLOOR<sup>4</sup> & DIGGER JACKSON<sup>5</sup>**

<sup>1</sup> Cork Ecology, Long Strand, Castlefreke, Clonakilty, Co. Cork, Ireland

<sup>2</sup> Bureau Waardenburg, Postbus 365, 4100 AJ Culemborg, the Netherlands

<sup>3</sup> Mainstream Renewable Power (Mainstream), Abbey Business Centre, Glasgow G2 5SG, UK

<sup>4</sup> Pelagica, Cotehill Farm, Collieston, Aberdeenshire AB41 8RX, UK

<sup>5</sup> Natural Research Projects Ltd, Brathens Business Park, Hill of Brathens, Banchory, Aberdeenshire AB31 4BY, UK

\*Email: [colin@corkeecology.net](mailto:colin@corkeecology.net)

**Introduction**

The large year-to-year changes that can occur in the marine environment make it difficult to detect and measure changes to bird populations caused by offshore wind farms.

Consequently, a robust survey design is essential to inform both pre-construction Environmental Impact Assessment (EIA) and subsequent post-construction monitoring of effects for projects that are consented.

In 2009 Mainstream Renewable Power Ltd was awarded the exclusive rights to develop the proposed 450-MW Neart na Gaoithe Wind Farm. The project is located in the outer Firth of Forth, 15.5 km off the Fife coast in eastern Scotland. The offshore Planning Application and Environmental Statement (ES) for Neart na Gaoithe Wind Farm was submitted to Marine Scotland in July 2012. Details of the Neart na Gaoithe project can be found at [www.neartnagaoithe.com](http://www.neartnagaoithe.com).

The use of before-after-control-impact (BACI) study designs to monitor the impact of offshore wind farms has become standard practice (e.g. Drewitt and Langston 2006, Fox *et al.* 2006, SNH 2009). However, there are concerns that BACI studies have little power to detect change, particularly those relating to potential displacement of seabirds (Underwood *et al.* 1991, 1994). In addition, for far-ranging species like seabirds there are inevitable difficulties with identifying truly independent control sites in the vicinity of the development site. Our study overcomes the shortcomings of BACI and instead uses a before-after-gradient (BAG) design (Mainstream Renewable Power 2009).

The BAG design assumes that:

- impacts decline with increasing distance from the source (Ellis & Schneider 1997, Morrison *et al.* 2008);
- any impacts due to displacement and habitat loss will be detected on the basis of changes in the distribution of seabirds in the area surveyed.

**Survey method**

The Study Area has two components: the Wind Farm Site and the surrounding 8-km-wide Buffer Area. Three years of monthly boat-based European Seabirds at Sea (ESAS) surveys were conducted between November 2009 and November 2012. The survey methods used followed COWRIE guidelines (Camphuysen *et al.* 2004) and all surveyors were ESAS accredited. The surveys covered the whole Study Area evenly, using parallel transects 2 km apart.

This paper forms part of the proceedings from the BOU conference *Marine Renewables and Birds*  
Other papers from these proceedings can be viewed at [www.BOUPROC.net](http://www.BOUPROC.net).

## Data analyses

Distance 6.0 software was used to estimate monthly numbers and densities for the most frequently recorded species. Species distribution was mapped at a resolution of 2 km across the Study Area.

## Results

Baseline surveys in Year 1 and Year 2 recorded:

- a total of 29 seabird species in the Study Area in Year 1, with 26 species in Year 2;
- Gannet *Morus bassanus*, Kittiwake *Rissa tridactyla*, Guillemot *Uria aalge*, Razorbill *Alca torda* and Puffin *Fratercula arctica* were the five most frequently recorded species, accounting for 83.4% all birds recorded in the Wind Farm Site in Year 1, and 91.4% in Year 2 (based on raw numbers).

### Between-year differences in abundance

The extent of between-year differences in abundance varied across species. For example, estimated numbers of Gannets between April and October for both the Wind Farm Site and the Buffer Area were higher in Year 2 than in Year 1 (Fig. 1). Estimated numbers of Gannets between November and March were more similar.





Figure 1. Monthly estimated Gannet population of Neart na Gaoithe Development & Buffer Areas in Years 1 and 2.


### Between-year variation in distribution

*This paper forms part of the proceedings from the BOU conference **Marine Renewables and Birds**  
Other papers from these proceedings can be viewed at [www.BOUPROC.net](http://www.BOUPROC.net).*

The extent of between-year differences in distribution varied across species and seasons. Particularly large differences were noted for auk species during the post-breeding season (September and October), when auks occurred in large numbers, and these are illustrated below.



**Figure 2.** Guillemot density in the Neart na Gaoithe Study Area in September and October, Years 1 and 2.



**Figure 3.** Puffin density in the Neart na Gaoithe Study Area in September and October, Years 1 and 2.

Between September and October of Year 1, Guillemots were generally evenly spread at high densities across the Study Area apart from in the southeast of the Buffer Area (Fig 2). In contrast, in the same period of Year 2, Guillemots were scarce



*This paper forms part of the proceedings from the BOU conference **Marine Renewables and Birds**  
Other papers from these proceedings can be viewed at [www.BOUPROC.net](http://www.BOUPROC.net).*

throughout most of the eastern part of the Study Area including much of the Wind Farm Site where they had occurred in high densities the previous year.

A similar difference in distribution pattern was observed for Razorbills in the post-breeding season, with fewer birds in the Wind Farm Site in September and October of Year 2 than in the same period in Year 1.

However, a very different pattern of distributional change was apparent for Puffins between the two post-breeding seasons. Between September and October of Year 1, highest densities of Puffins occurred in the eastern half of the Study Area (in contrast to Guillemot and Razorbill) (Fig. 3). Over the same period in Year 2, Puffins were generally evenly spread at moderate to high densities across the whole Study Area.

## Discussion

- The surveys provide adequate data to inform EIA and demonstrate that the Wind Farm Site and surrounding waters have high value for some seabird species.
- As expected, there is moderate to large annual variation in abundance and distribution in the baseline survey data, in both the Wind Farm Site and the surrounding 8 -km buffer.
- Different food availability conditions between years caused by hydrodynamic features and weather are likely to be the explanation for the variability shown here (within and outside the Wind Farm Site).
- Bringing proxies correlating with ecological differences between years into the analysis are therefore necessary (Macleane *et al.* 2012), to increase the power of detecting potential negative effects or to correct for other factors besides the wind farm (potentially yielding to false negative effects at the wind farm site as shown here).
- Also other factors such as trends in breeding numbers and year-to-year differences in breeding performances of the different species should be taken into account when considering differences in densities and distribution patterns at sea.
- The large (8 km wide) buffer area surveyed around the Wind Farm Site will enable the most statistically powerful BAG analyses available to be undertaken in future and, with correction for other factors, allow the effects of the wind farm on bird populations to be determined.

The application and ES for the offshore elements of the Neart na Gaoithe Wind Farm are currently being considered by Marine Scotland. A decision is expected in 2013. If consent is given for the development a construction and post-construction monitoring programme will be designed and the BAG approach applied. The Neart na Gaoithe team believe this will provide one of the best examples of a construction and post-construction monitoring plan so far implemented for offshore wind farms in the UK.

## References

- Camphuysen, C.J., Fox, T., Leopold, M.F. & Petersen, I.K. 2004. *Towards standardised seabirds at sea census techniques in connection with environmental impact assessments for offshore wind farms in the UK*. A report for COWRIE.
- Drewitt, A. & Langston, R.H.W. 2006. Assessing the impacts of wind farms on birds. *Ibis* 148: 29–42.  
<http://onlinelibrary.wiley.com/doi/10.1111/j.1474-919X.2006.00516.x/abstract>



*This paper forms part of the proceedings from the BOU conference **Marine Renewables and Birds**  
Other papers from these proceedings can be viewed at [www.BOUPROC.net](http://www.BOUPROC.net).*

- Ellis, J.I. & Schneider D.C. 1997. Evaluation of a gradient sampling design for environmental impact assessment. *Environmental Monitoring and Assessment* 48: 157–172.
- Fox, A.D., Desholm, M., Kahlert, J., Christensen, T.K. & Krag Petersen, I.B. 2006. Information needs to support environmental impact assessments of the effects of European marine offshore wind farms on birds. *Ibis* 148: 129–144.  
<http://onlinelibrary.wiley.com/doi/10.1111/j.1474-919X.2006.00510.x/abstract>
- Mainstream Renewable Power. 2009. *Neart na Gaoithe Proposed Offshore Wind Farm Scoping Report – Appendices*. Available at [www.neartnagaoithe.com](http://www.neartnagaoithe.com)
- Maclean, I.M.D., Rehfisch, M.M., Skov, H. & Thaxter C.B. 2012. Evaluating the statistical power of detecting changes in the abundance of seabirds at sea. *Ibis* 155: 113–126. <http://onlinelibrary.wiley.com/doi/10.1111/j.1474-919X.2012.01272.x/abstract>
- Morrison, M.L., Block, W.M., Strickland, M.D., Collier, B.A. & Peterson, M.J. 2008. *Wildlife Study Design*, 2nd edn. Berlin: Springer.
- SNH (Scottish Natural Heritage) 2009. *Guidance on Methods for Monitoring Bird Populations at Onshore Wind Farms*. Inverness: SNH.
- Underwood, A.J. 1991. Beyond BACI: experimental designs for detecting human environmental impacts on temporal variations in natural populations. *Aust. J. Mar. Freshwater Res* 42: 569–587.
- Underwood, A.J. 1994. On Beyond BACI: sampling designs that might reliably detect environmental disturbances. *Ecological Applications* 4: 3–15.